



Dr. Ramin Behmand | 204 23<sup>rd</sup> Avenue North, Nashville, TN 37203 | 615-576-3232

## Breast Augmentation Pre and Post-Operative Instructions

### General Information

- If you are a smoker (cigarettes or vaping) or use nicotine supplements, gum, or patches, it is very important that you stop at least 4 weeks prior to surgery. Smoking or using nicotine products can prevent blood flow that is vital for recovery after any surgery.
- It is best to stop any vitamin/medication such as Aspirin, NSAIDS (ibuprofen), Fish Oil, Vitamin E, ADD/ADHD medications and any diet pills/supplements- 1 week prior to surgery and 2 weeks after surgery. These will thin your blood. Tylenol is ok to take if needed.
- Make sure you have your prescriptions filled and waiting for you at home prior to surgery.
- It is important to have someone with you at home for the first 24-48hrs after surgery. You may feel groggy from the anesthesia and the pain medications.
- Do not eat or drink after midnight the day before surgery
- Wear something comfortable-a shirt that either buttons or zips in the front and pants that are easy to put on.
- Do not take pain medications on an empty stomach. Eat a light snack 30 minutes prior to taking pain medications. If you find you are nauseous after taking the pain medication, take your nausea medication 30 mins prior to your next pill and this should help.
- Try to stay on top of your pain at first after surgery. Taking your medication as prescribed at first will make for an easier first week after surgery.
- Do not drive or operate heavy machinery while taking pain medication. If you are still taking narcotics by the time of your first post-op appointment, then have someone drive you.
- Constipation is common after surgery from the pain medication and the anesthesia. It is safe to take Mira Lax or Colace after surgery to help prevent this. If you have no improvement after 7 days, please notify the office.
- Stay hydrated and move around as tolerated after surgery. Light walking is ok. Be careful when you change positions from sitting to standing as this may cause lightheadedness. Staying hydrated and moving as tolerated helps prevent blood clots.
- Take frequent deep breaths, pump your feet and ankles intermittently, as tolerated, while sitting/laying to help prevent blood clots.
- No strenuous activities, heavy lifting, straining or activities that would elevate your heart rate after surgery until cleared to do so by Dr. Behmand

### What to Expect

- After surgery you will be in a gauze bra. You will leave this on until you come in for your first post-op appointment.
- Pain, bruising, tightness and swelling can be expected- these typically peak around the 3<sup>rd</sup> day after surgery. The armpit area and sides of breast will be sore and hurt after surgery.

- After surgery your breast implants will look a little high on your chest and like a square box. This is normal, the implants will begin to fall into the pocket created for them and round out like a natural breast. We will let you know when you can move into a sports bra. No bra with underwire until 6 weeks post op.
- You will need to sleep in a reclining position for the first few weeks after surgery. This will help with swelling. Sleeping on 2 pillows is adequate.
- Do not push, pull, lift, climb, massage the surgical site, or do any strenuous activity, as this can elevate your blood pressure which may increase bleeding/bruising. In addition, these activities can lead to wound separation and prolong your recovery time. Dr. Behmand will let you know when it's safe to resume normal activity. Be mindful of how you reach with your arms and move after surgery.
- You will return to the office for your first post-op appointment typically 3-6 days after surgery.
- You may shower the next day after surgery. Keep your back to the water, do not let water beat down on the surgical site. Gently pat the area dry or let it air dry before putting your gauze bra back on. You may wash the gauze bra by hand if needed, and dry on the cold setting in your dryer.

**When to call our office:**

- If you develop severe shortness of breath, chest pain or are experiencing a medical emergency, call 911 and then notify your physician
- If you develop a fever of 101.4 or higher
- If your pain medication stops working
- If you have markedly increased pain, bruising, swelling or bloody drainage
- If you develop leg swelling or pain
- If you have persistent nausea or vomiting
- If you are uncertain about a medication or treatment
- If you are experiencing any adverse symptoms or changes that you do not understand
- If you have ANY questions or concerns

**Dr. Behmand's Office Phone Number: 615-576-3232**

If it is after office hours when you call, push one when prompted and you will be transferred to the answering service who will notify the person on call to call you back. Please do not email with medical questions, medical concerns or during a medical emergency as our email is not monitored after business hours or when we are seeing patients.

Signature and Date \_\_\_\_\_