



GRASSLAND
PLASTIC
SURGERY
SPECIALISTS

Dr. Ramin Behmand | 1200 Old Hillsboro Rd Franklin, TN 37069 | 615-576-3232

Blepharoplasty Pre/Post-Operative Instructions

General Information

- If you are a smoker (cigarettes or vaping) or use nicotine supplements, gum, or patches, it is very important that you stop at least 4 weeks prior to surgery. Smoking or using nicotine products can prevent blood flow that is vital for recovery after any surgery.
- It is best to stop any vitamin/medication such as Aspirin, NSAIDS (ibuprofen), Fish Oil, Vitamin E, ADD/ADHD medications and any diet pills/supplements- 1 week prior to surgery and 2 weeks after surgery. These will thin your blood. Tylenol is ok to take if needed.
- GLP1 drugs such as Ozempic, Wegovy, Mounjaro or Zepbound should also be stopped at least one week prior to surgery and may be resumed one week after.
- Make sure you have your prescriptions filled and waiting for you at home prior to surgery.
- It is important to have someone with you at home for the first 24-48hrs after surgery. You may feel groggy from the anesthesia and the pain medications.
- Do not eat or drink after midnight the day before surgery
- Wear something comfortable-a shirt that either buttons or zips in the front and pants that are easy to put on.
- Do not take pain medications on an empty stomach. Eat a light snack 30 minutes prior to taking pain medications. If you find you are nauseous after taking the pain medication, take your nausea medication 30 mins prior to your next pill and this should help.
- Try to stay on top of your pain at first after surgery. Taking your medication as prescribed at first will make for an easier first week after surgery.
- Do not drive or operate heavy machinery while taking pain medication. If you are still taking narcotics by the time of your first post-op appointment, then have someone drive you.
- Constipation is common after surgery from the pain medication and the anesthesia. It is safe to take Mira Lax or Colace after surgery to help prevent this. If you have no improvement after 7 days, please notify the office.
- Stay hydrated and move around as tolerated after surgery. Light walking is ok. Be careful when you change positions from sitting to standing as this may cause lightheadedness. Staying hydrated and moving as tolerated helps prevent blood clots.
- No strenuous activities, heavy lifting, straining or activities that would elevate your heart rate after surgery until cleared to do so by Dr. Behmand

What to Expect

- Swelling, bruising and discoloration are normal. Swelling usually peaks around the 3rd day and then goes down from there.
- You may apply ice compresses gently to the eye area to help with swelling. You can apply this for 20 minutes, rest 20 minutes and repeat as often as you like. This will help with swelling. This really helps the first week after surgery.
- Its normal to have blurry vision after surgery. Sometimes it feels like you may have something in your eye as well and feel scratchy this is normal.
- Use liquid tears eye drops to keep your eyes moist as needed. Get these from the pharmacy prior to your surgery.
- Do not rub your eyes.
- After surgery you will need to keep your head elevated. Do not bend your head upside down, bend over and try not to flex the neck too hard. Be mindful of how you turn your neck and your head movements after surgery, turning your head gently and slowly.
- You will need to sleep with your head elevated after surgery. You may place 2 pillows behind your back or sleep in a recliner. Some patient's find a travel pillow comfortable. Sleeping elevated helps with swelling.
- You will return to the office 2-5 days after surgery for your first post op appointment
- You may shower after surgery. Let the water hit you from behind and use a washcloth to wash your face, staying away from the eyelids. You will want to be very careful with your face as it is still healing. Do not wash the eye area until cleared by Dr. Behmand to do so.

When to call our office:

- If you develop severe shortness of breath, chest pain or are experiencing a medical emergency, call 911 and then notify your physician
- If you develop a fever of 101.4 or higher
- If your pain medication stops working
- If you have markedly increased pain, bruising, swelling or bloody drainage
- If you develop leg swelling or pain
- If you have persistent nausea or vomiting
- If you are uncertain about a medication or treatment
- If you are experiencing any adverse symptoms or changes that you do not understand
- If you have ANY questions or concerns

Dr. Behmand's Office Phone Number: 615-576-3232

If it is after office hours when you call, push one when prompted and you will be transferred to the answering service who will notify the person on call to call you back. Please do not email with medical questions, medical concerns or during a medical emergency as our email is not monitored after business hours or when we are seeing patients.

Signature and Date_____